

GENERAL INDEX TO VOLUME XXIV

New scientific names of plants and the final members of new combinations are printed in **bold face type**; synonyms and page numbers having reference to figures and plates, in *italics*; and previously published names and all other matter, in ordinary type.

A

- Acanthaceae, Panamanian, determined by E. C. Leonard, 205
 Achimenes longiflora, 202
 Actinochaete, 1, 310; *flicina*, 2, 6, 8, 10; *flicina*, 263, 310, 311, 466, 468, 470, 474, 480
 Ailanthus altissima, 442, 499; *Cacodendron*, 499; *glandulosa*, 264, 264, 499; *peregrina*, 264, 499; *peregrina*, 264; *procera*, 499
 Albionia *Peregrina*, 264, 264, 499
 Allomarkgrafia *Brenesiana*, 15; *ovalis*, 15; *plumeriaefolia*, 16; *subtubulosa*, 16
 Alloplectus ichthyoderma, 205; *rubida*, 204
 Aloe sp., development of the male gametophyte in, 161, 163, 164
 Amyris *tozifera*, 306
 Anacardiaceae: A new genus of, 1; Studies in the I, 263, II, 499
 Anacardium, 1
 Aphelandra *Seibertii*, 205, 207
 Aplopappus, 228, 248, 250; *ciliatus*, 250; *occidentalis*, 250
 Apocynaceae, New or otherwise noteworthy, of Tropical America, V, 11
 Appunia guatemalensis, 209; *Seibertii*, 208
 Araliaceae, Panamanian, determined by P. C. Standley, 196
 Ardisia *Seibertii*, 198
 Aster, 225; *glutinosus*, 225; *spathularis*, 225; *spathulatus*, 225
 Astereae, 248
 Aurelia, 227; *amplexicaulis*, 228; *decurrens*, 228

B

- Bailey, L. H., Panamanian plants determined by: *Palmae*, 180; *Rosaceae*, 189
 Barkley, Fred A. A new genus of Anacardiaceae, 1; A monographic study of *Rhus* and its immediate allies in ANN. MO. BOT. GARD., VOL. 24, 1937.

- North and Central America, including the West Indies, 265; Studies in the Anacardiaceae, I, 263, II, 499
 Berry, Edward C. A preliminary list of the lichens of central Missouri, with ecological notes, 211
 Betula *triphylla*, 394
 Blake, S. F., Panamanian plants determined by: *Compositae*, 210; *Polygalaceae*, 190
 Blastomycotic dermatitis, 584
 Bletia *campanulata*, 185
 Bromeliaceae, Panamanian, determined by L. B. Smith, 180
 Bursera *bipinnata*, 3, 311

C

- Calanthe mexicana, 184
 Canal Zone, Missouri Botanical Garden Tropical Station at Balboa, 175, 176
 Canna latifolia, 182
 Carcinogens, 583
 Casselia, 27; *denticulata*, 140; *lanceolata*, 89; *maritima*, 34; *paniculata*, 43; *parviflora*, 34, *virginica*, 70
 Castellania *tropicalis*, 584
 Catopsis *nutans*, 180
 Cavendishia *Smithii*, 197; *Wercklei*, 197
 Centropogon *diocleus*, 209
 Cerbera *Ahouai*, 11; *nitida*, 11
 Cerinthodes, 27; *alpinum*, 107; *lanceolatum*, 89; *maritimum*, 35; *oblongifolium*, 123; *paniculatum*, 43; *virginicum*, 70
 Chamaedorea *costaricana*, 180
 Chert and flint rocks as a habitat of lichens, 213
 Chiriquí: Collections in the provinces of, Coclé, and Panamá, by R. J. Seibert during the summer of 1935, 177, 178; scene near the headwaters of the Rio Chiriqui Viejo, 177; scene at Cerro Punta, 177; view of Llanos del Volcan, 178
 Chrysophthalmum, 228; *andinum*, 229
 Cissus *Martiniana*, 191; *salutaris*, 191

- Coclé: Collections in the provinces of Chiriquí, Panamá, and, by R. J. Seibert during the summer of 1935, 177, 179; Llano Bonito, views of 178, 179
- Coffea, 537; oleifolia, 538; paniculata, 538
- Collections in the provinces of Chiriquí, Coclé, and Panamá, by R. J. Seibert, during the summer of 1935, 177
- Compositae, Panamanian, determined by S. F. Blake, 210
- Conopholis americana, 202
- Contributions toward a Flora of Panamá, based upon collections by expeditions from the Missouri Botanical Garden Tropical Station, Balboa, C. Z., 175; I, Collections in the provinces of Chiriquí, Coclé, and Panamá, by R. J. Seibert during the summer of 1935, 177
- Cotinus, 1, 300; americanus, 301, 466, 468, 470, 474, 478; Coggyria, 300, 301, 478; Copallinus, 321; cotinoides, 301; Metopium, 308
- Cotinus, 303, 314
- Cyperaceae, Panamanian, determined by Fr. Hugh O'Neil, 180

D

- Daphnopsis Seibertii, 192
- Datisca hirta, 326
- Demetria, 226; glutinosa, 227; spathulata, 227
- Diastema exiguum, 202, var. Ilacinum, 203
- Dieliptera, 544
- Dodge, Carroll W. and Bertha Sanford. Some effects of methyl cholanthrene on the morphology and growth of yeasts, 583
- Donia, 226; squarrosa, 226
- Doniana, 228
- Doronicum, 226; glutinosum, 226
- Dryopteris macradenia, 180

E

- Ecological notes on the lichens of central Missouri, 211
- Elleanthus capitatus, 182
- Epidendrum Parkinsonianum, 184; pygmaeum, 184
- Eucharis Bouchei, 181

F

- Fagara, 263
- Faramaea, 537; caerulea, 537; Guayaquilensis, 538; jasminoides, 537; latifolia, 538; Montevicensis, 538; odoratissima, 538; sessilifolia, 538; stipulacea, 538

- Flora of Panamá, Contributions toward a, 175

- Fuchsia arborescens, 196; Hemsleyana, 194, 196; mixta, 194; pulchella, 195; Seleriana, 194, 196

- Funastrum clausum, 200; Seibertii, 199

G

- Gametophyte, the male, observations on the development of, in certain monocots, 161, 174
- Gases, effect of, on growth of lichens, 212
- Gasteria, development of male gametophyte in, 162, 163
- Gesneriaceae, Panamanian, determined by C. V. Morton, 202
- Gilibertia arborea, 197; stenodonta, 196
- Granules, refractive, in gametophyte of Tradescantia, 167; in Aloe, 169
- Grindelia, the genus: cytological aspects of, 260; geographical distribution of, 229, 232; history of, 225; phylogeny of, 252, chart showing, 254; relationships of, 247, chart showing, 251; specific concept of, 261; Studies in, III, 225
- Grindelia acutifolia, 256; aggregata, 230; angustifolia, 228; aphanactis, 230, 255; arenicola, 230, 242, 259; arizonica, 246, 253, var. microphylla, 256, var. stenophylla, 256; aphanactis, 230; Blakei, 230; camporum, 230, 239, 242, 258, var. Davyi, 230, 239, 242, 258, var. parviflora, 258; columbiana, 240; decumbens, 230, 256, var. subincisa, 256; foliosa, 228; fruticosa, 228; glutinosa, 228; grandiflora, 230, 245, 260; Greenmanii, 260; Hallii, 257; Havardii, 246, 253; hirsutula, 259, var. brevisquama, 261; humilis, 230; integrifolia, 238, 260, var. virgata, 238, 260; inuloides, 225, 228; laciniata, 256; lanceolata, 230, 245; littoralis, 233, 256; microcephala, 260, var. adenodonta, 261; nana, 230, 239, 242, var. altissima, 242, 258, var. turbinella, 258; Nelsonii, 260; oolepis, 233; oxylepis, 253, var. eligulata, 253, 255; perennis, 229, 243, 255; procera, 230, 257; pulchella, 228; resinosa, 228; revoluta, 256; rubricaulis, 231, 259, 261, var. bracteosa, 261, var. elata, 231, 259, 261, var. latifolia, 231, var. platyphylla, 236, 259, f. pilosa, 261, var. robusta, 231, 259, 261; scabra, 246, 256, var. neomexicana, 246, 256; speciosa, 228; squarrosa, 228, 229, 243, 255, var. nuda, 231, 255, var. serrulata, 229, 255; stricta,

- 228, 236, 259, var. *aestuarina*, 236, var. *Andersonii*, 236, var. *collina*, 231, 236, var. *Hendersonii*, 231, var. *lanata*, 236, 238, var. *macrophylla*, 236; *subalpina*, 256; *subdecurrens*, 253; *tenuella*, 260; *texana*, 230, 245; *Volkensii*, 228
- Grindeliaceae*, 248
- Guzmania polycephala*, 180
- H
- Habenaria setifera*, 182
- Hackelia virginiana*, 141
- Happ, George B. Monograph of *Tetramerium* and *Henrya*, 501
- Heliantheae*, 248
- Helosis mexicana*, 188
- Henrya*, 544; Monograph of *Tetramerium* and, 501
- Henrya*, 544; *Augustiniana*, 567; *Barclayana*, 561, 582; *brevifolia*, 547; *Conzattii*, 560; *costata*, 548; *costata*, 551, var. *glandulosa*, 546; *Donnell-Smithii*, 563; *flava*, 537, 550; *grandifolia*, 561; *gualanensis*, 537, 553, 582; *imbricans*, 558; *insularis*, 546, 578, 582; *laxa*, 557; *longipes*, 549; *mephitica*, 562; *Ortegana*, 552; *pilosa*, 556; *puberula*, 559; *reticulata*, 566; *rupicola*, 564, 582; *scorpioides*, 538, 546, 554, 576, 582, var. *latifolia*, 556; *scorpioides*, 546; *Silvestrii*, 567; *yucatanensis*, 551
- Henryastrum*, 567; *Augustinianum*, 567; *Silvestrii*, 567
- Hippoglossum*, 27; *maritimum*, 35; *virginicum*, 70
- Hoorebekia*, 228; *chiloensis*, 228
- Huntleya Burtii*, 185
- Hypericum fastigiatum*, 192
- I
- Inula*, 225; *glutinosa*, 226; *serrata*, 226
- Ipomoea muricata*, 201, forma *alba*, 201
- J
- Johnson, G. T. and Ruth E. Peck. Observations on the male gametophyte in certain monocots, 161
- Justicia racemulosa*, 538
- K
- Killip, E. P., Panamanian *Passifloraceae* determined by, 192
- L
- Leonard, E. C., Panamanian *Acanthaceae* determined by, 205
- Lichens of central Missouri, A preliminary list of the, with ecological notes, 211
- Limestone ledges as a habitat of lichens, 214
- Liparis Wendlandii*, 183
- Lipocarpha Sellowiana*, 180
- Lithospermum*, 27; *corymbosum*, 141; *denticulatum*, 140; *Drummondii*, 140; *marginatum*, 89; *maritimum*, 34; *paniculatum*, 43; *pulchrum*, 70; *pumilum*, 34
- Lithraea*, 414; *laurina*, 415
- Llano Bonito, Panama, view of, looking toward Pacific Ocean, 178; view across "badlands" of, 179
- Llanos del Volcan, Chiriqui, 178
- Lobadium*, 314, 341, 390; *amentaceum*, 394; *aromaticum*, 395; *suaveolens*, 394; *trifoliatum*, 395
- Lobeliaceae*, Panamanian, determined by F. E. Wimmer, 209
- Loranthaceae*, Panamanian, determined by William Trelease, 187
- Lycaste Dowiana*, 185
- M
- Malaxis brachyrrhyncha*, 182; *excavata*, 182; *macrostachya*, 182; *Parthonii* var. *denticulata*, 182
- Malosma*, 1, 414; *laurina*, 415, 466, 468, 470, 474, 496
- Maxon, W. R. Panamanian *Polypodiaceae* determined by, 180
- Melastomaceae*, Panamanian, determined by P. C. Standley, 193
- Mertensia*, A monograph of the genus, in North America, 17
- Mertensia*, 26; *alaskana*, 41, 48; *alba*, 55; *alpina*, 106, 107; *alpina* var. *brevistyla*, 102, var. *humilis*, 105, var. *perplexa*, 110; *ambigua*, 87; *amoena*, 130; *amplifolia*, 98; *arizonica*, 62, 63, var. *Grahami*, 65, var. *Leonardi*, 64, var. *subnuda*, 66, var. *umbratilis*, 64; *Bakeri*, 118, 118, var. *Osterhoutii*, 120; *Bakeri* var. *amoena*, 130, var. *lateriflora*, 118, var. *subglabra*, 123; *bella*, 40, 40; *brachycalyx*, 49; *brachyloba*, 97; *brevistyla*, 102, 103; *brevistyla* var. *obtusiloba*, 107; *caelestina*, 114; *californica*, 84; *campanulata*, 68, 69; *cana*, 115; *canescens*, 115; *ciliata*, 74, 75, var. *latiloba*, 84, var. *stomatocoides*, 84, var. *subpubescens*, 83; *ciliata* var. *longipedunculata*, 74, var. *polyphylla*, 75, var. *punctata*, 75, f. *candida*, 83; *Clokeyi*, 95; *congesta*, 100; *Cooperae*, 130; *coriacea*, 113, var. *dilatata*, 113; *coronata*, 125; *corym-*

- bosa, 141; *Cusickii*, 130; *cynoglossoides*, 114; *denticulata*, 140; *Drummondii*, 140; *Eastwoodiae*, 41; sect. *Eumertensia*, 29; *Fendleri*, 97, var. *pubens*, 98; *foliosa*, 125, var. *amoena*, 130, f. *Cusickii*, 130, var. *nevadensis*, 125, var. *pubescens*, 130, var. *subcalva*, 123, f. *Macbridei*, 123; *franciscana*, 55; *fusiformis*, 100, *grandis*, 55; *Horneri*, 136; *humilis*, 105, 105; *incongruens*, 75; *infirma*, 87; *intermedia*, 125; *laevigata*, 49, var. *brachycalyx*, 49; *lanceolata*, 88, 89, var. *brachyloba*, 97, var. *Fendleri*, 97, var. *pubens*, 98, var. *secundorum*, 95; *lan- ceolata* var. *aptera*, 88, var. *lineariloba*, 110, var. *myosotifolia*, 118, var. *viridis*, 110; *lateriflora*, 118; *Leonardi*, 64; *leptophylla*, 49; *lineariloba*, 110; *linearis*, 88; *longiflora*, 136, 136; *longiflora* var. *Horneri*, 136, var. *pulchella*, 136; *MacDougalii*, 121, 122; *marginata*, 88; *maritima*, 34, 35; *maritima* f. *albiflora*, 34; *media*, 95; *membranacea*, 42; *mexicana*, 67; *micrantha*, 95; *muriculata*, 114; *myosotifolia*, 118; *Nelsonii*, 123, 125; sect. *Neuranthis*, 29; *nevadensis*, 125; *nivalis*, 118; *nutans*, 125, subsp. *subcalva*, 123; *oblongifolia*, 123, 123, var. *amoena*, 130, var. *nevadensis*, 125; *oblongifolia* var. *nimbata*, 130; *obtusiloba*, 106; *oreophila*, 134, 135; *ovata*, 110; *pallida*, 75; *Palmeri*, 43; *paniculata*, 42, 44, var. *alaskana*, 48, var. *borealis*, 49; *paniculata* var. *laevigata*, 49, var. *longisepala*, 42, var. *nivalis*, 118, var. *platyphylla*, 59, var. *subcordata*, 60, f. *leptophylla*, 49; *papillosa*, 88, var. *fusiformis*, 100, var. *lineariloba*, 110; *Parryi*, 110; *parviflora*, 34; *perpleza*, 110; *picta*, 74; *pilosa*, 140; *platyphylla*, 59, var. *subcordata*, 60; *polyphylla*, 74; *praecox*, 125; *pratensis*, 55, f. *alba*, 55, var. *borealis*, 49; *pubescens*, 130; *pulchella*, 136, subsp. *glauca*, 136; *pulmonarioides*, 70; *punctata*, 74; *refracta*, 118; *rivularis*, 42; *Sampsonii*, 64; *secundorum*, 95; *sibirica* var. *Drummondii*, 140; *siskiyouensis*, 40; sect. *Steenhammery*, 28; sect. *Stenhammery*, 28; *stenoloba*, 125; *stomatocoides*, 84; *subcordata*, 60; *subpubescens*, 83; *symphytoides*, 125; *toyabensis*, 61; *toyabensis* var. *subnuda*, 66; *tubiflora*, 125; *Twedyi*, 106; *umbratilis*, 87; *virginica*, 69, 70; *viridis*, 110, var. *caelestina*, 114, var. *cana*, 115, var. *cynoglossoides*, 114, var. *dilatata*, 113, var. *parvifolia*, 115; *viridula*, 110.
- Metastelma glaberrimum*, 200; *pedunculare*, 201
- Methyl cholanthere, Some effects of, on the morphology and growth of yeasts, 583
- Metopium*, 1, 303; *Brownei*, 304, 308; *Brownei* var. *brachycarpum*, 308; *Linnaei*, 306, 308, var. *Oxymetopium*, 306; *Metopium*, 306; *toxisferum*, 263, 306, 466, 468, 470, 474; *venosum*, 305, 480
- Missouri Botanical Garden Tropical Station, Balboa, C. Z., Administration Building, 176; collections by expeditions from, 175
- Missouri, central, A preliminary list of the lichens of, with ecological notes, 211
- Mitosis in *Tradescantia* gametophyte, 165
- Moisture, effect of, on growth of lichens, 212
- Monocots, Observations on the development of the male gametophyte in certain, 161, 174
- Monograph of *Tetramerium* and *Henrya*, 501
- Monograph of the genus *Mertensia* in North America, 17
- Monographic study of *Rhus* and its immediate allies in North and Central America, including the West Indies, 265
- "Monte Lirio," as a collecting base, in Chiriquí, Panama, 178
- Morton, C. V., Panamanian Gesneriaceae determined by, 202
- Mouriria brunneicalyx*, 193; *parviflora*, 194
- Mycocandida onychophila*, 584; effect of methyl cholanthere on growth of, 588
- Myrica trifoliata*, 395
- Myrsinaceae, Panamanian, determined by P. C. Standley, 198
- N
- Napeanthus*, 203; *repens*, 203
- Neostyphonia*, 314, 341, 355; *integri- folia*, 360; *ovata*, 365
- O
- Ocotea Whitel*, 188
- O'Neill, Fr. Hugh, Panamanian Cyperaceae determined by, 180
- Orchidaceae, Panamanian, determined by C. Schweinfurth, 182

Orchids at Missouri Botanical Garden
Tropical Station, 175
Oxalis (*Ionoxalis*) *chiriquensis*, 190

P

Pachycormus *discolor*, 442
Palmae, Panamanian, determined by
L. H. Bailey, 180
Panama: Collections in the provinces of
Chiriquí, Coeló, and, by R. J. Seibert
during the summer of 1935, 177; Con-
tributions toward a flora of, 175, I,
177
Passiflora *apetala*, 192
Passifloraceae, Panamanian, determined
by E. P. Killip, 192
Peck, Ruth E., G. T. Johnson and. Ob-
servations on the development of the
male gametophyte in certain mono-
cots, 161
Peperomia *antoni*, 185; *liratinerve*,
186; *marmorum*, 186; *rivi-vetusti*,
186; *Seibertii*, 185; *subrepens*, 187;
valliculæ, 186
Pernettya *coriacea*, 197
Persea *alpigena*, 306
Philostemon, 417; *radicans*, 425
Philostemum, 417;
Phinsea *caripensis*, 204; *lacerata*, 203
Phoradendron *corynarthron* var. *Sei-
bertii*, 187
Picramnia *polyantha*, 442
Pierasma *antillana*, 441
Pigment, yellow, of lichens, effect of
sunlight on, 212
Piperaceae, Panamanian, determined by
William Trelease, 185
Platynema, 27; *paniculata*, 43
Pleurothallis *cardiothallis*, 184; *phyllo-
cardia*, 184; *triangulabia*, 183;
Tuerckheimii, 184
Plumeriopsis, 11; *Ahouai*, 11
Pneumaria, 27; *maritima*, 34; *virginica*,
70
Pocophorum, 417
Polygala *leptocaulis*, 190
Polygalaceae, Panamanian, determined
by S. F. Blake, 190
Polymnia *maculata*, 210
Polypodiaceae, Panamanian, determined
by W. R. Maxon, 180
Polypodium *astrolepis*, 180; *dissimile*,
180
Pongelion *glandulosum*, 499
Powell, C. W., gift of orchid garden of,
to Missouri Botanical Garden, 175
Pseudosmodingium *perniciosum*, 263,
442; *Pterocarpus*, 263; *Pterocarpus*,
500; *rhodifolium*, 442, 500

Pulmonaria, 27; *alpina*, 107; *canaden-
sis*, 70; *ciliata*, 75; *denticulata*, 140;
elegans, 43; *elliptica*, 141; *glabra*, 70;
lanceolata, 88; *marginata*, 88; *mari-
tima*, 34; *oblongifolia*, 123; *panicu-
lata*, 43; *parviflora*, 34; *pilosa*, 140;
virginica, 70

Q

Quercus *insignis*, 187; *eugeniaefolia*,
187; *Seemanni*, 187

R

Rauwolfia *indecora*, 12
Rhipidopteris *flabellulata*, 180
Rhododendron *catawbiense*, 441
Rhocidium, 314, 341, 386; *cinereum*,
387; *glabellum*, 387; *microphyllum*,
386; *Potosinum*, 387; *retusum*, 387;
rugulosum, 387; *vestitum*, 387
Rhoec, development of the male gameto-
phyte in, 162
Rhus, 312; *acutiloba*, 426; *albida*, 331;
allophylloides, 392, 494; *americana*,
301, 330; *Andrieuxii*, 379, 466, 470,
474; *angustiarum*, 332; *antillana*, 441;
aprica, 332; *arboorea*, 441; *arbores-
cens*, 441; *arbuscula*, 332; *arguta*,
332; *aromatica*, 342, 390, 394, 468,
472; *aromatica*, 435, var. *glabra*, 394,
var. *mollis*, 409, var. *schmidelioides*,
393, var. *trilobata*, 399; *Arsenel*, 346,
476, 482; *Ashei*, 340; *asplenifolia*,
332; *atrovirens*, 332; *auriculata*, 332;
bahamensis, 426; *Barclayi*, 350, 468;
bipinnata, 339; *Blodgettii*, 426; *bore-
alis*, 340; *Cacodendron*, 499; *calo-
phylla*, 332; *Canadense*, 331; *cana-
dense*, 394; *canadensis*, 326, var. *il-
linoensis*, 394, var. *serotina*, 406, var.
simplicifolia, 410; *Canadensis* var.
mollis, 409, var. *quinata*, 413; *caro-
liniana*, 340; *Carolinianum*, 326; *caro-
linense*, 331; *catawbiense*, 441; *Cav-
anillesii*, 441; *chondroloma*, 370;
choriophylla, 378; *ciliolata*, 384; *cis-
montana*, 332; *coccinea*, 331; *Copal-
lina*, 316; *Copallina* var., 321, var.
lanceolata, 322, var. *leucantha*, 321;
copallina var. *angustifolia*, 322, f.
integrifolia, 323, f. *serrata*, 323, var.
arborescens, 316, f. *crispa*, 317, var.
extensa, 316, var. *latifolia* f. *angusti-
alata*, 321, f. *latialata*, 316, var. *neso-
phila*, 316, var. *salicifolia*, 316; *coral-
lina*, 316; *Coriaria*, 1, 315; *costari-
censis*, 351; *cotinoides*, 301; *Cotinus*,
300, 301; *crenata*, 394; *crenatifolia*,

441; *divaricata*, 433; *diversifolia*, 423; *diversiloba*, 423, f. *quinquifolia*, 423, f. *radicans*, 423; *Duckeri*, 371, 438; *elegans*, 331, var. *glauca*, 331, var. *superba*, 331; *elegantula*, 332; *Emoryi*, 409; *eximia*, 435; sect. *Eutoxiocodendron*, 419; *flicifolia*, 329; *Filicina*, 1, 2, 310, 311; *flicina*, 2, 3; *floridana*, 427; *Florita*, 441; *frutescens*, 326; *fruticosum*, 442; *Galeottii*, 348; *glabra*, 1, 331, 466, 468, 470, 472, 474, var. *borealis*, 340, 472, var. *laciniata*, 399; *glabra*, var. *cismontana*, 331, var. *coccinea*, 331, var. *dioica*, 331, var. *elegans*, 331, var. *hermaphrodita*, 331, f. *laciniata*, 339, var. *occidentalis*, 331, var. *Sandbergii*, 340; *glabra* × *typhina*, 330; *glabrum* var. *canadense*, 331, var. *carolinense*, 331; *gracilis*, 326; *Greeni*, 433; *gymnoclada*, 330; *Hapemanii*, 332; *Hartmanii*, 344, 433; *heterophyllum*, 442; *Hindsiana*, 360; *hirta*, 326, var. *dissecta*, 329, var. *laciniata*, 329, var. *typhina*, 326; *humile*, 426; *humilis*, 426; *hybrida*, 330; *hypselerodendrum*, 331; *ilinoensis*, 394, var. *formosa*, 394; *integrifolia*, 356, 360, 466, 470, var. *cedrosensis*, 363, 436; *integrifolia*, 365, var. *serrata*, 360; *ithacensis*, 332; *jalscana*, 345; *juglandifolia*, 436, var. *Lindeniana*, 436, var. *Samo*, 436; *Kearneyi*, 363, 434; *laevicaulis*, 331; *lanceolata*, 322; *laurina*, 415; *Lentii*, 357; *lenticifolia*, 317; *leucantha*, 321; *Lentii*, 357; *Lindeniana*, 437; *lineata*, 442; *lineatifolia*, 442; *littoralis*, 434; sect. *Lobadium*, 390; *lobata*, 423; *longula*, 332; *lucida*, 441; *ludoviciana*, 332; *macrophylla*, 442; *macropoda*, 333, 492; *macrothyrsa*, 331; *media*, 332; *methopium*, 308; *metopia*, 308; *metopioidea*, 441; *Metopium*, 308, 348; *metopium*, 306; *Michauxii*, 324; *microcarpa*, 426; *microphylla*, 386, 387, 466, 468, 470, 494, var. *vestita*, 387; *mollis*, 358; *Muelleri*, 359, 472, 484; *Nelsonii*, 375, 490; *nitens*, 332; *Nortonii*, 394; *oaxacana*, 376, 472; *obtusifolia*, 321; *occidentalis*, 331; *oreophila*, 332; *Osterhoutii*, 399; *ovata*, 365, 468, 476, var. *Traskiae*, 368, 486; *ozyacanthoides*, 400; *Oxymetopium*, 306; *pachyrrhachia*, 373; *pachyrrhachia*, 375; *Palmeri*, 351; *peregrina*, 264, 442, 499; *perniciosa*, 442; *petiolata*, 332; *pistachiaefolia*, 317; *polyantha*, 442; *potentillaefolia*, 1, 3, 311; pro-

fusa, 385, 492; sect. *Pseudoschmaltzia*, 369; sect. *Pseudosumac*, 343; *pteroearpa*, 500; *Pterocarpus*, 263, 442, 500; *pubescens*, 420, 427; *pulchella*, 332; *pulvinata*, 330; *pumilum*, 324; *pyramidata*, 332; *quercifolia*, 420; *quinquefolia*, 308; *radicans*, 425, var. *microcarpa*, 426, var. *volubilis*, 426, var. *vulgaris*, 426; *refescens*, 442; sect. *Rhoetidum*, 386; *rhomboides*, 435; *rubifolia*, 349, var. *subcordata*, 349; *Rydbergii*, 427; *sambucina*, 332; *Samo*, 437; *Sandbergii*, 340; *sanguinea*, 331; *saxatilis*, 442; *Scandens*, 426; *scandens*, 426; *Schiedeana*, 380; *Schiedeana*, 373, 375, f. *vestita*, 374; *schinoides*, 442; subg. *Schmaltzia*, 341; *schmidelioides*, 393, var. *potosinensis*, 394; *sempervirens*, 377, var. *pachyrrhachia*, 375; *Sinense*, 499; *sorbifolia*, 332; *Standleyi*, 358; *striata*, 436; sect. *Styphonla*, 355; *suavcolens*, 394; *subcordata*, 349; *Succedaneum*, 264, 264, 374, 442, 499; subg. *Sumac*, 315; *Tepetate*, 382, 490; *terebinthifolia*, 343, 353, 466, 470, var. *Barclayi*, 350, var. *Loeseneri*, 354, var. *pilosissima*, 351; *tessellata*, 332; *tetlatin*, 442; *tetlatsiam*, 3; *Tetlatsiam*, 1, 3, 311; *Tetlasian*, 1; *Toxicarium*, 426; *toxicarium*, 426; *toxicodendron*, 419, 420, 426, subsp. *diversiloba*, 423, var. *eximia*, 435, f. *malacotrichocarpum*, 434, var. *microcarpon*, 426, var. *quercifolium*, 420, var. *radicans*, 426, f. *radicans*, 426, var. *Rydbergii*, 426, var. *vulgare*, 426, var. *vulgaris*, f. *radicans*, 426, f. *volubilis*, 426; *Toxicodendron*, 426; sect. *Trichocarpae*, 315, 342, 390; *tridentatum*, 426; *trilobata*, 399, 466, var. *anisophylla*, 411, var. *arenaria*, 408, var. *glaberrima*, 399, var. *pilosissima*, 409, var. *quinata*, 413, var. *racemulosa*, 409, var. *serotina*, 406, 466, 470, 472, var. *simplicifolia*, 410; *typhina*, 326, 474, 476, var. *laciniata*, 329; *typhina* var. *arborescens*, 326, f. *dissecta*, 329, var. *flicifolia*, 329, var. *flicina*, 329, var. *frutescens*, 326, var. *viridiflora*, 326; *typhium*, 326; *Uillosum*, 426; *utahensis*, 410; *valida*, 332; *variclobata*, 423; *variifolium*, 442; *Veatchiana*, 442; *venenata*, 438; *venosa*, 305; *Vernicifera*, 263, 438; *Vernix*, 436, 438; *verrucosa*, 435; *vestita*, 374, var. *Ghiesbreghtii*, 375; *virens*, 2, 368, 377, 468, 472; *viridiflora*, 326, 331, var. *canadense*, 326

Rhus, 300, 303, 310, 390, 414, 417
Rhus-Toxicodendron, 418; *radicans*, 427;
toxicodendrum, 427; *Vernix*, 438
 Rio Chiriquí Viejo, 178; scene near the
 headwaters of, 177; Llanos del Vol-
 can, seen through the gorge of, 178
 Rosaceae, Panamanian, determined by
 L. H. Bailey, 189
 Rubiaceae, Panamanian, determined by
 P. C. Standley, 208
Rubus glaucus, 189

S

Sabouraud glucose agar, yeast cultures
 grown on, 584
Saccharomyces ellipsoideus, 584; effect
 of methyl cholanthrene on growth of,
 588, on morphology of, 586; cham-
 pagne strain, 584, 587, 590; Tokay
 strain, 584
 Sandstone ledges as habitat of lichens,
 213
Sansevieria, development of the male
 gametophyte in, 162, 163, 164
Sapindus marginatus, 441
Schmalzia, 314, 341, 390; *affinis*, 410;
anisophylla, 411; *anomala*, 409;
arenaria, 408; *aromatica*, 395; *Ashei*,
 340; *Bakeri*, 400; *botryoides*, 400;
cissodes, 410; *cognata*, 400; *copallina*,
 317; *crataegifolia*, 395; *crenata*, 395;
cruciata, 411; *elegantula*, 400; *Em-
 oryi*, 409; *formosa*, 395; *glabra*,
 332; *glabrata*, 400; *glauca*, 400;
glomerata, 400; *hederacea*, 411; *hirta*,
 326; *hirtella*, 400; *illinoensis*, 395;
lanceolata, 323; *lasiocarpa*, 406; *leio-
 carpa*, 400; *leucantha*, 321; *mala-
 cophylla*, 409; *Michauxii*, 324; *mi-
 crophylla*, 387; *Nortonii*, 395;
obtusifolia, 321; *Oregana*, 411; *oxy-
 acanthoides*, 400; *pubescens*, 400;
pulchella, 409; *puncticulata*, 411;
quercifolia, 400; *quinata*, 413; *race-
 mulosa*, 409; *ribifolia*, 410; *sabulosa*,
 400; *scaberula*, 409; *serotina*, 406;
serrata, 395; *simplicifolia*, 410; *stra-
 minea*, 411; *suaveolens*, 395; *sub-
 pinnata*, 400; *tridophylloides*, 400;
trilobata, 400; *trinervata*, 411; *virens*,
 377
Schmalzia, 314, 341
Schmidelia Cominia, 441
 Schweinfurth, C., Panamanian Orchi-
 daceae determined by, 182
 Seibert, Russell J., Robert E. Woodson,
 Jr. and. Collections in the provinces of
 Chiriquí, Coelá, and Panamá, by R. J.

Seibert, during the summer of 1935,
 177
 Smith, L. B., Panamanian Bromeliaceae
 determined by, 180
 Soil, as a habitat of lichens: of glades
 and abandoned fields, 214; of dry,
 rocky, woody hills, 214; of moist
 woods, 214
 Solidagineae, 248
Spathelia rhoifolia, 500
Spiranthes Scopulariae, 182
 Standley, P. C., Panamanian plants de-
 termined by: Araliaceae, 196; Melas-
 tomaceae, 193; Myrsinaceae, 198;
 Rubiaceae, 208; Thymelaeaceae, 192
Steenhamera virginica, 70
Steenhamera, 27, 29; *maritima*, 34;
virginica, 70
Stellaria ovata, 188
 Steyermark, Julian A. Studies in
 Grindelia. III, 225
 Studies in Grindelia. III, 225
 Studies in the Anacardiaceae. I, 263,
 II, 499
Styphonia, 314, 341, 355; *integrifolia*,
 356, 360; *mollis*, 358; *serrata*, 360
 Sunlight, effect of, on growth of lichens,
 211

T

Terebinthus Brovvnii, 308; *Brownii*,
 305, 308
 Tetrad in certain monocots, 162, 163,
 163, 164, 174
 Tetramerium, Monograph of, and Hen-
 rya, 501
Tetramerium, 501, 503, 508; *aureum*,
 533, 582; *Calderonii*, 519; *coeruleum*,
 537; *costatum*, 548; *diffusum*, 534;
 sect. *Eutetramerium*, 509; *flavum*,
 537, 550; *fruticosum*, 517; *genicu-
 latum*, 537; *glandulosum*, 532, 582;
glutinosum, 522; *gualanense*, 537,
 553; *Hilli*, 515; *Hintonii*, 524; *his-
 pidum*, 511, 527, 572, 580, 582, var.
Greenmanii, 531; *jasmynoides*, 537;
Langlassei, 523; *latifolium*, 538; *lep-
 tocaule*, 516; *macrostachyum*, 520;
montevideense, 538; *multiflorum*, 538;
nemorum, 521; *nervosum*, 511, 570,
 578, 580, 582, var. *acuminatum*, 568,
 var. *angustifolium*, 512; *nervosum*
 var. *hispidum*, 527; *occidentale*, 538;
odoratissimum, 538; *oleaeifolium*, 538;
ovalifolium, 518; *ovatum*, 513; *pa-
 niculatum*, 538; *platystegium*, 535,
 574, 582; *polystachyum*, 525; *racemu-
 losum*, 538; *rubrum*, 536; *scabrum*,
 532; *scorpioides*, 538, 546, 554, 561;

- sessilifolium*, 522; *sessilifolium*, 538;
Standleyi, 514; *stipulaceum*, 538;
tenuissimum, 526; sect. *Torreyella*,
 509, 535
Tetramerium, 508, 537, 544
Thalictrum podocarpum, 188
Thevetia Ahouai, 11; *nitida*, 11
Thouinia acuminata, 263
Thuraria, 226
 Thymelaeaceae, Panamanian, deter-
 mined by P. C. Standley, 192
Tonduzia macrantha, 12
Toxicodendron, 1, 417; *aboriginum*, 434;
Altissimum, 499; *Andrieuxii*, 379;
Arboreum, 441; *arizonicum*, 425;
bitermum, 435; *Blodgettii*, 425;
comarophyllum, 422; *compactum*, 420;
copallinum 317, var. *latifolium*, 317;
coriaceum, 422; *Crenatum*, 342, 390,
 395; *cuneatum*, 395; *desertorum*, 425;
divaricatum, 433; *diversiloba*, 422;
dryophilum, 423; *ezimium*, 435;
fothergilloides, 425; *Glabrum*, 425;
glabrum, 332; *goniocarpum*, 425;
hesperium, 425; *Hindianum*, 360; *in-*
tegrifolium, 360; *isophyllum*, 423;
laetevirens, 425; *laurinum*, 415; *Len-*
tii, 357; *lobadioides*, 423; *longipes*,
 425; *macrocarpum*, 425; *macrophyll-*
um, 442; *magnum*, 425; *microphyll-*
um, 387; *molle*, 358; *monticola*,
 420; *Negundo*, 425; *oxycarpum*, 423;
phaseoloides, 425; *Pinnatis*, 438; *Pin-*
natum, 438; *potentillifolium*, 3, 311;
pubescens, 420, 425; *pumilum*, 324,
 425; *punctatum*, 425; *quercifolia*, 420,
 498; *radicans*, 419, 425, 498, var. *di-*
varicata, 433, var. *eximia*, 435, var.
littoralis, 434, var. *verrucosa*, 435;
radicans var. *microcarpa*, 425, var.
normale, 425, var. *volubile*, 425; *rhom-*
boideum, 435; *rubifolium*, 349; *rufes-*
cens, 425; *Rydbergii*, 425; *Schiede-*
anum, 381; *sempervirens*, 377; *Ser-*
ratum, 425; *striata*, 436, 496; *tere-*
binthifolium, 353; *Toxicodendron*,
 420; *trilobatum*, 400; *triphyllum*,
 425, var. *trilobatum* f. *glaberrimum*,
 400; *typhinum*, 326; *vaccarium*, 423;
vaciarum, 423; *verniciifera*, 263;
 sect. *Vernix*, 436; *Vernix*, 436, 438,
 466, 468, 470, 474; *Vernix*, 438; *ver-*
rucosum, 435; *Volubile*, 425; *Vulgare*,
 425
Toxicodendron, 310, 314, 390, 419
Tradescantia: *hirsutiflora*, 174, and *T.*
paludosa, 167, development of male
 gametophyte in, 161; unnamed hy-
 brida, development of male gameto-
 phyte in, 161, 174
 Tree trunks, as a habitat of lichens, 215
 Trelease, William, Panamanian plants
 determined by: *Loranthaceae*, 187;
Piperaceae, 185
 Tropical America, collecting plants in,
 177
Turpinia, 314, 341, 390; *aromatica*, 395;
glabra, 395; *pubescens*, 395; *suaveo-*
lens, 395
 V
 Vacuoles in *Tradescantia* gametophyte,
 165, 171, 174
 Valle de Anton, 179, view into, from
 southern rim, 179
Vallesia antillana, 13; *Balleyana*, 14;
flexuosa, 14; *glabra*, 13; *montana*, 15
Vernix, 418, 436
Vincetoxicum chiriquense, 199
Vriesia subsecunda, 181
 W
 Water courses, beds of, as a habitat of
 lichens, 215
Weinmannia pinnata, 189
 Wimmer, F. E., Panamanian *Lobelia-*
ceae determined by, 209
 Williams, Louis O. A monograph of the
 genus *Mertensia* in North America, 17
Winkleria, 27
 Wood, old and decaying, as a habitat of
 lichens, 215
 Woodson, Robert E. Jr. New or other-
 wise noteworthy *Apocynaceae* of trop-
 ical America, V, 11; and Russell J.
 Seibert. Contributions toward a Flora
 of Panama. I. Collections in the prov-
 inces of Chiriquí, Coelá, and Panamá,
 by R. J. Seibert during the summer of
 1935, 175
 Y
 Yeasts, Some effects of methyl cholanth-
 rene on the morphology and growth
 of, 583
 Z
Zymonema dermatitidis, 584

